



DRUG MARKET WITHDRAWAL
DESMOPRESSIN ACETATE TABLETS 0.1 MG AND
DESMOPRESSIN ACETATE TABLETS 0.2MG
0.1 MG (NDC 68462-275-01) & 0.2 MG (NDC 68462-276-01)
(Glenmark)

July 26, 2023

Dear Customer,

Glenmark is voluntarily requesting a Market Withdrawal at wholesaler level involving the following Desmopressin Acetate Tablets batches;

DESMOPRESSIN ACETATE TABLETS 0.1 MG

S. No.	NDC	Batch No.	Expiry Date
1.	68462-275-01	20210715	10/2023
2.	68462-275-01	20220094	01/2024
3.	68462-275-01	20220155	02/2024

DESMOPRESSIN ACETATE TABLETS 0.2 MG

S. No.	NDC	Batch No.	Expiry Date
1.	68462-276-01	20210651	09/2023
2.	68462-276-01	20210664	09/2023
3.	68462-276-01	20210666	09/2023
4.	68462-276-01	20210716	10/2023
5.	68462-276-01	20210818	11/2023
6.	68462-276-01	20220051	12/2023
7.	68462-276-01	20220058	12/2023
8.	68462-276-01	20220060	12/2023
9.	68462-276-01	20220100	01/2024



S. No.	NDC	Batch No.	Expiry Date
10.	68462-276-01	20220132	02/2024
11.	68462-276-01	20220133	02/2024
12.	68462-276-01	20220137	02/2024

Market withdrawal of these batches have been initiated due to Out Of Specification (OOS) was reported in test of Content Uniformity (By HPLC) for the product "Desmopressin Acetate Tablets 0.2 mg" B. No. 20210314 A.R. No. ST3230680 for condition 25°C/60% RH during testing of Shelf life interval where one tablet failed during CU testing. The subjected batch # 20210314 has been expired in the month of May 2023 and hence not included in this market withdrawal.

No assignable root cause could be identified for the observed out of specification result during laboratory and manufacturing investigation. Glenmark is initiating market withdrawal all of these batches voluntarily as FDA decided that this is not an official recall but a market withdrawal because none of these batches subject to Glenmark's voluntary market withdrawal failed at release or on stability to date. Therefore, out of an abundance of caution, Glenmark is requesting market withdrawal for these batches.

See details of product batches listed in above table on page 1 & 2 and refer enclosed product labels for ease in identifying the product.

Examine your inventory and if you have any batches specified in table of page 1 & 2 then quarantine it immediately.

We are requesting the batches specified in the above table to be returned to Qualanex, LLC (address below) using the Postage Paid Product Return label that was provided in your Market Withdrawal Return Packet.

Qualanex, LLC
1410 Harris Road
Libertyville, IL 60048

Please complete and return the enclosed response form preferably within 72 hours of receipt of this notification. Please either fax your response to 847-737-3719 or email to recall@qualanex.com.



If you have any questions regarding your market withdrawal return, please contact Qualanex at 800-505-9291.

Thank you for your cooperation,

Sincerely,

GLENMARK PHARMACEUTICALS INC., USA

Thomas Callaghan Digitally signed by Thomas Callaghan
Date: 2023.07.26 12:18:18 -04'00'

Thomas Callaghan

Executive Director - Regulatory Affairs, North America

US Agent for Glenmark Pharmaceuticals Limited

Tel.: (201) 675-9447

Fax: (201) 205-1378

E-mail: thomas.callaghan@glenmarkpharma.com

Enclosure(s):

Product Labels

Market Withdrawal Return Response Form

MARKET WITHDRAWAL RETURN RESPONSE FORM

DESMOPRESSIN ACETATE TABLETS 0.1 MG AND DESMOPRESSIN ACETATE TABLETS 0.2MG- 0.1 MG (NDC 68462-275-01) & 0.2 MG (NDC 68462-276-01)

(Qualanex Market Withdrawal # 392)

DESMOPRESSIN ACETATE TABLETS 0.1 MG

S. No.	NDC	Batch No.	Expiry Date
1.	68462-275-01	20210715	10/2023
2.	68462-275-01	20220094	01/2024
3.	68462-275-01	20220155	02/2024

DESMOPRESSIN ACETATE TABLETS 0.2 MG

S. No.	NDC	Batch No.	Expiry Date
1.	68462-276-01	20210651	09/2023
2.	68462-276-01	20210664	09/2023
3.	68462-276-01	20210666	09/2023
4.	68462-276-01	20210716	10/2023
5.	68462-276-01	20210818	11/2023
6.	68462-276-01	20220051	12/2023
7.	68462-276-01	20220058	12/2023
8.	68462-276-01	20220060	12/2023
9.	68462-276-01	20220100	01/2024
10.	68462-276-01	20220132	02/2024
11.	68462-276-01	20220133	02/2024
12.	68462-276-01	20220137	02/2024

Please fill out and either fax to (847) 737-3719 or email your completed response form to recall@qualanex.com.

NDC: _____ Lot: _____

Quantity Being Returned: Full/Sealed Bottles Partial/Open Bottles

MARKET WITHDRAWAL RETURN RESPONSE FORM

(Qualanex Market Withdrawal # 392)

NDC: _____ Lot: _____

Quantity Being Returned:	Full/Sealed Bottles	Partial/Open Bottles
	_____	_____

NDC: _____ Lot: _____

Quantity Being Returned:	Full/Sealed Bottles	Partial/Open Bottles
	_____	_____

We have examined our inventories and have none of the above market withdrawal product in stock:

(Initials) _____

Form Completed By: (Please Print)

Name/Title: _____

Phone: _____

E-Mail: _____ Fax # _____

Company Name: _____

Company DEA: _____

Company Address: _____

City, State, Zip _____

Signature: _____


Wholesaler: _____

Wholesaler Acct. No. _____

Date: _____

I have read and understand the market withdrawal instructions provided in the Market Withdrawal Letter received on: _____ [Date]

JAR SIZE : 80 CC
 SAME SIZE ARTWORK
 LABEL SIZE 105 X 45 MM




MINIMUM FONT SIZE: 4 PT

GLENMARK PHARMACEUTICALS LTD.		DATE:	PANTONE SHADE NO: <input type="checkbox"/> BLACK <input type="checkbox"/> 186 C
PRODUCT NAME: Desmopressin Acetate Tab 0.1 mg		PKG. DEV.:	Item code, Version, Consistency of Design, overprint area, Pack size, Dimensions & Layout
ITEM CODE: PE40140 VERSION: 0217-1		RA	Regulatory Text
PHARMACODE:		QA:	Entire Text
COUNTRY: USA		PRODUCTION:	Machine Suitability
LOCATION: COLVALE - GOA		REMARKS:	
PACK: LABEL (100 TABLETS)			
ACTUAL SIZE: 105 mm x 45 mm			
SPECIFICATION: A UV VARNISH COATED FASPRINT NG/PERMANENT (HITAC)/ PET 23 STICKER LABEL FROM AVERY DENISON PRINTED AS PER APPROVED ARTWORK. BATCH PRINTING AREA REMAINS UNVARNISHED			

Handwritten notes:
 Radhan
 2/5/17
 3/5/17
 10/5/17
 10/5/17


JAR SIZE : 60 CC
 SAME SIZE ARTWORK
 LABEL SIZE 105 X 45 MM



MINIMUM FONT SIZE: 4 PT

GLENMARK PHARMACEUTICALS LTD.	DATE:	PANTONE SHADE NO: <input type="checkbox"/> BLACK <input type="checkbox"/> 2945 C	
		<input type="checkbox"/> 186 C	
PRODUCT NAME: Desmopressin Acetate Tab 0.2 mg	PKG. DEV.:	Item code, Version, Consistency of Design, overprint area, Pack size, Dimensions & Layout	
ITEM CODE: PE40141 VERSION: 0217-1	RA	Regulatory Text	
PHARMACODE:	QA:	Entire Text	
COUNTRY: USA	PRODUCTION:	Machine Suitability	
LOCATION: COLVALE - GOA	REMARKS:		
PACK: LABEL (100 TABLETS)			
ACTUAL SIZE: 105 mm x 45 mm			
SPECIFICATION: A UV VARNISH COATED FASPRINT NG/PERMANENT (HITAC)/ PET 23 STICKER LABEL FROM AVERY DENISON PRINTED AS PER APPROVED ARTWORK. BATCH PRINTING AREA REMAINS UNVARNISHED			

Handwritten notes:
 Bradhan 2/17/17
 3/5/17
 5/17



Desmopressin Acetate Tablets, 0.1mg and 0.2mg

Rx Only

DESCRIPTION

Desmopressin Acetate Tablets are a synthetic analogs of the natural pituitary hormone D-arginine vasopressin (AVP), an antidiuretic hormone affecting renal water conservation. It is chemically defined as follows:

Mol. Wt. 1129.27 Empirical Formula: $C_{20}H_{31}N_5O_4 \cdot CH_3COO$



1-(3-mercaptopropionic acid)-D-arginine vasopressin monoacetate (salt).

Desmopressin acetate tablets, for oral administration, contain either 0.1 or 0.2 mg desmopressin acetate, USP. Inactive ingredients include: lactose monohydrate, potato starch, talc, magnesium stearate and povidone.

CLINICAL PHARMACOLOGY

Desmopressin Acetate Tablets, contain as active substance, desmopressin acetate, a synthetic analog of the natural hormone arginine vasopressin.

Control Diabetes Insipidus:

Dose response studies in patients with diabetes insipidus have demonstrated that oral doses of 0.025 mg to 0.4 mg produced clinically significant antidiuretic effects. In most patients, doses of 0.1 mg to 0.2 mg produced optimal antidiuretic effects lasting up to eight hours. With doses of 0.4 mg, antidiuretic effects were observed for up to 12 hours; measurements beyond 12 hours were not recorded. Increasing oral doses produced dose dependent increases in the plasma levels of desmopressin acetate.

The plasma half-life of desmopressin acetate followed a monoexponential time course with a $t_{1/2}$ value of 1.5 to 2.5 hours which was independent of dose.

The bioavailability of Desmopressin Acetate oral tablets is about 5% compared to intranasal desmopressin acetate, and about 0.19% compared to intravenous desmopressin acetate. The time to reach maximum plasma desmopressin acetate levels ranged from 0.9 to 1.5 hours following oral or intranasal administration, respectively. Following administration of Desmopressin Acetate Tablets, the onset of antidiuretic effect occurs at around 1 hour, and it reaches a maximum at about 4 to 7 hours based on the measurement of increased urine osmolality.

The use of Desmopressin Acetate Tablets in patients with an established diagnosis will result in a reduction in urinary output with an accompanying increase in urine osmolality. These effects usually will allow resumption of a more normal life style, with a decrease in urinary frequency and nocturia.

There are reports of an occasional change in response to the intranasal formulations of Desmopressin Acetate (Desmopressin Acetate Nasal Spray and Desmopressin Acetate Rhinal Tube). Usually, the change occurred over a period of time greater than six months. This change may be due to decreased responsiveness, or to shortened duration of effect. There is no evidence that this effect is due to the development of binding antibodies, but may be due to a local inactivation of the peptide. No loss of effect was observed in the 46 patients who were treated with Desmopressin Acetate Tablets for 12 to 44 months and no serum antibodies to desmopressin were detected.

The change in structure of arginine vasopressin to desmopressin acetate resulted in less vasopressor activity and decreased action on visceral smooth muscle relative to enhanced antidiuretic activity. Consequently, clinically effective antidiuretic doses are usually below the threshold for effects on vascular or visceral smooth muscle. In the four long-term studies of Desmopressin Acetate Tablets, no increases in blood pressure in 46 patients receiving Desmopressin Acetate Tablets for periods of 12 to 44 months were reported.

In one study, the pharmacodynamic characteristics of Desmopressin Acetate Tablets and intranasal formulation were compared during an 8-hour dosing interval at steady state. The doses administered to 36 hydrated (water loaded) healthy male adult volunteers every 6 hours were 0.1, 0.2, 0.4 mg orally and 0.01 mg intranasally by rhinal tube. The results are shown in the following table:

Treatment	Total Urine Volume in ml	Maximum Urine Osmolality in mOsm/kg
0.1 mg PO q6h	-3989.3 (149.5)	514.3 (21.9)
0.2 mg PO q6h	-4229.9 (149.5)	686.3 (21.9)
0.4 mg PO q6h	-4698.8 (149.5)	769.3 (21.9)
0.01 mg IN q6h	-4844.9 (149.5)	754.1 (21.9)

(SE) = Standard error of the mean

With respect to the mean values of total urine volume decrease and maximum urine osmolality increase from baseline, the 90% confidence limits estimated that the 0.4 mg and 0.2 mg oral doses produced between 55% and 110% and 84% to 95% of pharmacodynamic activity, respectively, when compared to the 0.01 mg intranasal dose.

While both the 0.2 mg and 0.4 mg oral doses are considered pharmacodynamically similar to the 0.01 mg intranasal dose, the pharmacodynamic data on an inter-subject basis was highly variable and, therefore, individual dosing is recommended.

In another study in diabetes insipidus patients, the pharmacodynamic characteristics of Desmopressin Acetate Tablets and intranasal formulations were compared over a 12-hour period. Ten fluid-controlled patients under age 18 were administered tablet doses of 0.2 mg and 0.4 mg, and intranasal doses of 0.01 mg and 0.02 mg.

Treatment	Urine Volume in mL/min	Maximum Urine Osmolality in mOsm/kg
0.01 mg IN	0.3 (0.15)	717.0 (224.63)
0.02 mg IN	0.3 (0.25)	761.8 (298.82)
0.2 mg PO	0.3 (0.12)	878.3 (147.91)
0.4 mg PO	0.2 (0.15)	787.2 (73.34)

(SD) = Standard Deviation

All four dose formulations (0.01 mg IN, 0.02 mg IN, 0.2 mg PO and 0.4 mg PO) have a similar, pronounced pharmacodynamic effect on urine volume and urine osmolality. At two hours after study drug administration, mean urine volume was 4 mL/min and urine osmolality was >500 mOsm/kg. Mean plasma osmolality remained

relatively constant over the time course recorded (0 to 12 hours). A statistical separation from baseline did not occur at any dose or time point. In these patients, the 0.2 mg tablets and the 0.01 mg intranasal spray exhibited similar pharmacodynamic profiles as did the 0.4 mg tablets and the 0.02 mg intranasal spray formulation. In another study of adult diabetes insipidus patients previously controlled on Desmopressin Acetate intranasal spray, after one week of self-titration from spray to tablets, patients' diuresis was controlled with 0.1 mg Desmopressin Acetate Tablets three times a day.

Primary Nocturnal Enuresis:

Two double-blind, randomized, placebo-controlled studies were conducted in 340 patients with primary nocturnal enuresis. Patients were 5-17 years old, and 72% were males. A total of 329 patients were evaluated for efficacy. Patients were evaluated over a two-week baseline period in which the average number of wet nights was 10 (range 4-14). Patients were then randomized to receive 0.2, 0.4, or 0.8 mg of Desmopressin Acetate or placebo. The pooled results after two weeks are shown in the following table:

	Placebo (n = 85)	0.2 mg/day (n = 78)	0.4 mg/day (n = 82)	0.8 mg/day (n = 85)
Baseline	10 (0.3)	11 (0.3)	10 (0.3)	10 (0.3)
Reduction from Baseline	1 (0.3)	3 (0.4)	3 (0.4)	4 (0.4)
Percent Reduction from Baseline	10%	27%	30%	40%
p-value vs placebo	—	<0.05	<0.05	<0.05

Patients treated with Desmopressin Acetate Tablets showed a statistically significant reduction in the number of wet nights compared to placebo-treated patients. A greater response was observed with increasing doses up to 0.8 mg.

In a six month, open-label extension study, patients completing the placebo-controlled studies were started on 0.2 mg/day desmopressin acetate, and the dose was progressively increased until the optimal response was achieved (maximum dose 0.8 mg/day). A total of 230 patients were evaluated for efficacy; the average number of wet nights/2 weeks during the untreated baseline period was 10 (range 4-14), and the average duration (SD) of treatment was 4.2 (1.8) months. Twenty-five (25) patients (11%) achieved a complete or near complete response (≤ 2 wet nights/2 weeks) and did not require titration to the 0.8 mg/day dose. The majority of patients (198 of 230, 86%) were titrated to the highest dose. When all dose groups were combined, 128 (56%) showed at least a 50% reduction from baseline in the number of wet nights/2 weeks, while 87 (38%) patients achieved a complete or near complete response.

Human Pharmacokinetics:

Desmopressin Acetate is mainly excreted in the urine. A pharmacokinetic study conducted in healthy volunteers and patients with mild, moderate, and severe renal impairment (n=24, 6 subjects in each group) receiving single dose desmopressin acetate (2mg) injection demonstrated a difference in desmopressin acetate terminal half-life. Terminal half-life significantly increased from 3 hours in normal healthy patients to 9 hours in patients with severe renal impairment. (See CONTRAINDICATIONS).

INDICATIONS AND USAGE

Control Diabetes Insipidus:

Desmopressin Acetate Tablets are indicated as antidiuretic replacement therapy in the management of central diabetes insipidus and for the management of the temporary polyuria and polydipsia following head trauma or surgery in the pituitary region. Desmopressin Acetate tablets are ineffective for the treatment of nephrogenic diabetes insipidus.

Patients were selected for therapy based on the diagnosis by means of the water deprivation test, the hypertonic saline infusion test, and/or response to antidiuretic hormones. Continued response to Desmopressin Acetate tablets can be monitored by measuring urine volume and osmolality.

Primary Nocturnal Enuresis:

Desmopressin Acetate Tablets are indicated for the management of primary nocturnal enuresis. Desmopressin Acetate Tablets may be used alone or in an adjunct to behavioral conditioning or other non-pharmacologic interventions.

CONTRAINDICATIONS

Desmopressin Acetate Tablets are contraindicated in individuals with known hypersensitivity to desmopressin acetate or to any of the components of Desmopressin Acetate Tablets.

Desmopressin Acetate is contraindicated in patients with moderate to severe renal impairment (defined as a creatinine clearance below 30 mL/min).

Desmopressin Acetate is contraindicated in patients with hyponatremia or a history of hyponatremia.

WARNINGS

1. Very rare cases of hyponatremia have been reported from world-wide postmarketing experience in patients treated with desmopressin acetate. Desmopressin Acetate is a potent antidiuretic which, when administered, may lead to water intoxication and/or hyponatremia. Unless properly diagnosed and treated hyponatremia can be fatal. Therefore, fluid restriction is recommended and should be decreased with the patient and/or guardian. Careful medical supervision is required.

2. When Desmopressin Acetate Tablets are administered, in particular in pediatric and geriatric patients, fluid intake should be adjusted downward to decrease the potential occurrence of water intoxication and hyponatremia. (See PRECAUTIONS, Pediatric Use and Geriatric Use.) All patients receiving desmopressin acetate therapy should be observed for the following signs or symptoms associated with hyponatremia: headache, nausea/vomiting, decreased serum sodium, weight gain, restlessness, fatigue, lethargy, disorientation, depressed reflexes, loss of appetite, irritability, muscle weakness, muscle spasms or cramps and abnormal mental status such as hallucinations, decreased consciousness and confusion. Severe symptoms may include one or a combination of the following: seizures, coma and/or respiratory arrest. Particular attention should be paid to the possibility of the rare occurrence of an extreme decrease in plasma osmolality that may result in seizures which could lead to coma.

3. Desmopressin Acetate should be used with caution in patients with habitual or psychogenic polydipsia who may be more likely to drink excessive amounts of water, putting them at greater risk of hyponatremia.

PRECAUTIONS

General:

Intranasal formulations of desmopressin acetate at high doses and Desmopressin Acetate Injection have infrequently produced a slight elevation of blood pressure which disappears with a reduction of dosage. Although this effect has not been observed when single oral doses up to 0.8 mg have been administered, the drug should be used with caution in patients with coronary artery insufficiency and/or hypertensive cardiovascular disease, because of a possible rise in blood pressure. Desmopressin Acetate should be used with caution in patients with conditions associated with fluid and electrolyte imbalance, such as cyclic fibrils, heart failure and renal disorders because these patients are prone to hyponatremia.

GLENMARK GENERICS LTD.		DATE: 04/03/2015	PANTONE SHADE NO: Black
PRODUCT NAME: LFLT DESMOPRESSIN ACETATE TAB US GPI-AE	PKG. DEV.:	Item code, Version, Consistency of Design, overprint area, Pack size, Dimension & Layer	<i>Diadhar 9/4/15</i>
ITEM CODE: FE32425 VERSION: 0315-1	RA:	Regulatory Text	<i>9/4/15</i>
PHARMACODE:	QA:	Entire Text	<i>9/4/15</i>
COUNTRY:	PRODUCTION:	Machina Suitability	<i>9/22/15</i>
LOCATION: Goa	REMARKS:		
PACK : 40 GSM BIBLE PAPER			
ACTUAL SIZE: 200X280 MM			
SPECIFICATION: 32X32 MM			

Rare severe allergic reactions have been reported with Desmopressin Acetate. Anaphylaxis has been reported rarely with intravenous and intranasal administration of desmopressin acetate but not with Desmopressin Acetate Tablets.

Laboratory Tests:

Central Diabetes Insipidus:
 Laboratory tests for monitoring the patient with central diabetes insipidus or post-surgical or head trauma-related polyuria and polydipsia include urine volume and osmolality. In some cases, measurements of plasma osmolality may be useful.

Drug Interactions:
 Although the pressor activity of desmopressin acetate is very low compared to its antidiuretic activity, large doses of Desmopressin Acetate Tablets should be used with other pressor agents only with careful patient monitoring. The concomitant administration of drugs that may increase the risk of water intoxication with hypotension, (e.g., tricyclic antidepressants, selective serotonin re-uptake inhibitors, chlorpromazine, opiate analgesics, NSAIDs, barbiturates and carbamazepine) should be performed with caution.

Carcinogenicity, Mutagenicity, Impairment of Fertility:
 Studies with desmopressin acetate have not been performed to evaluate carcinogenic potential, mutagenic potential or effects on fertility.

Prepregnancy:

Category B:
 Fertility studies have not been done. Teratology studies in rats and rabbits at doses from 0.05 to 10 mcg/kg/day (approximately 0.1 times the maximum systemic human exposure in rats and up to 38 times the maximum systemic human exposure in rabbits based on surface area, mg/m²) revealed no harm to the fetus due to Desmopressin Acetate Tablets. There are, however, no adequate and well-controlled studies in pregnant women. Because animal studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Several publications where desmopressin acetate was used in the management of diabetes insipidus during pregnancy are available; these include a few anecdotal reports of congenital anomalies and low birth weight babies. However, no causal connection between these events and desmopressin acetate has been established. A three year Swedish epidemiologic study of the use of desmopressin acetate in pregnant women with diabetes insipidus found the rate of birth defects to be no greater than that in the general population; however, the statistical power of this study is low. As opposed to preparations containing natural hormones, desmopressin acetate in antidiuretic doses has no uterotropic action and the physician will have to weigh the possible therapeutic advantages against the possible risks in each case.

Nursing Mothers:

There have been no controlled studies in nursing mothers. A single study in postpartum women demonstrated a minimal change in plasma, but little if any change in excretable desmopressin acetate in breast milk following an intranasal dose of 0.1 mg.

It is not known whether the drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when desmopressin acetate is administered to nursing mothers.

Pediatric Use:

Central Diabetes Insipidus:
 Desmopressin Acetate Tablets have been used safely in pediatric patients, age 4 years and older, with diabetes insipidus for periods up to 44 months. In younger pediatric patients the dose must be individually adjusted in order to prevent an excessive decrease in plasma osmolality leading to hyponatremia and possible convulsions; dosing should start at 0.05 mg (1/2 of the 0.1 mg tablet). Use of Desmopressin Acetate Tablets in pediatric patients requires careful fluid intake restrictions to prevent possible hyponatremia and water intoxication. Fluid restriction should be discussed with the patient and/or guardian. (See WARNINGS.)

Primary Nocturnal Enuresis:

Desmopressin Acetate Tablets have been safely used in pediatric patients age 6 years and older with primary nocturnal enuresis for up to 6 months. Some patients respond to a dose of 0.2 mg; however, increasing responses are seen at doses of 0.4 mg and 0.6 mg. No increase in the frequency or severity of adverse reactions or decrease in efficacy was seen with an increased dose or duration. The dose should be individually adjusted to achieve the best results. Treatment with desmopressin for primary nocturnal enuresis should be interrupted during acute intercurrent illness characterized by fluid and/or electrolyte imbalance (e.g., systemic infection, fever, recurrent vomiting or diarrhea) or under conditions of extremely hot weather, vigorous exercise or other conditions associated with increased water intake.

Geriatric Use:

Clinical studies of Desmopressin Acetate Tablets did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function. Desmopressin Acetate is contraindicated in patients with moderate to severe renal impairment (defined as a creatinine clearance below 50 mL/min). (See CLINICAL PHARMACOLOGY, Human Pharmacokinetics and CONTRAINDICATIONS.)

Use of Desmopressin Acetate Tablets in geriatric patients requires careful fluid intake restrictions to prevent possible hyponatremia and water intoxication. Fluid restriction should be discussed with the patient. (See WARNINGS.)

ADVERSE REACTIONS

Infrequently, large doses of the intranasal formulations of Desmopressin Acetate and Desmopressin Acetate Injection have produced transient headache, nausea, flushing and mild abdominal cramps. These symptoms have disappeared with reduction in dosage.

Central Diabetes Insipidus:

In long-term clinical studies in which patients with diabetes insipidus were followed for periods up to 44 months of Desmopressin Acetate Tablet therapy, transient increases in AST (SGOT) no higher than 1.5 times the upper limit of normal were occasionally observed. Elevated AST (SGOT) returned to the normal range despite continued use of Desmopressin Acetate Tablets.

Primary Nocturnal Enuresis:

The only adverse event occurring in 20% of patients in controlled clinical trials with Desmopressin Acetate Tablets that was probably, possibly, or remotely related to study drug was headache (4% desmopressin acetate, 9% placebo).

Other:

The following adverse events have been reported; however their relationship to desmopressin acetate has not been established: observed delirium, dizziness, and edema-weight gain. See WARNINGS for the possibility of

water intoxication and hyponatremia.

Pediatric Monitoring:

There have been case reports of hyponatremic convulsions associated with concurrent use with the following medications: cyclophosphamide and indinavir.

OVERDOSEAGE

Signs of overdose may include confusion, drowsiness, constraining headache, problems with passing urine and rapid weight gain due to fluid retention. (See WARNINGS.) In case of overdose, the dose should be reduced, frequency of administration decreased, or the drug withdrawn according to the severity of the condition. There is no known specific antidote for desmopressin acetate. The patient should be observed and treated with appropriate symptomatic therapy. An oral LD₅₀ has not been established. Oral doses up to 0.2 mg/kg/day have been administered to dogs and rats for 6 months without any significant drug-related toxicities reported. An intravenous dose of 2 mg/kg in mice demonstrated no effect.

DOSEAGE AND ADMINISTRATION

Central Diabetes Insipidus:
 The dosage of Desmopressin Acetate Tablets must be determined for each individual patient and adjusted according to the diurnal pattern of response. Response should be estimated by two parameters: adequate duration of sleep and adequate, not excessive, water turnover. Patients previously on intranasal Desmopressin Acetate therapy should begin tablet therapy twelve hours after the last intranasal dose. During the initial dose titration period, patients should be observed closely and appropriate safety parameters measured to assure adequate response. Patients should be monitored at regular intervals during the course of Desmopressin Acetate Tablet therapy to assure adequate antidiuretic response. Modifications in dosage regimen should be implemented as necessary to assure adequate water turnover. Fluid restriction should be observed. (See WARNINGS, PRECAUTIONS, Pediatric Use and Geriatric Use.)

Adults and Children:

It is recommended that patients be started on doses of 0.05 mg (1/2 of the 0.1 mg tablet) two times a day and individually adjusted to their optimum therapeutic dose. Most patients in clinical trials found that the optimal dosage range is 0.1 mg to 0.6 mg daily, administered in divided doses. Each dose should be separately adjusted for an adequate diurnal rhythm of water turnover. Total daily dosage should be increased or decreased in the range of 0.1 mg to 1.2 mg divided into two or three daily doses as needed to obtain adequate antidiuresis. See Pediatric Use subsection for special considerations when administering desmopressin acetate to pediatric diabetes insipidus patients.

Geriatric Use:

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function. (See CLINICAL PHARMACOLOGY, Human Pharmacokinetics, CONTRAINDICATIONS, and PRECAUTIONS, Geriatric Use.)

Primary Nocturnal Enuresis:

The dosage of Desmopressin Acetate Tablets must be determined for each individual patient and adjusted according to response. Patients previously on intranasal Desmopressin Acetate therapy can begin tablet therapy the night following (24 hours after) the last intranasal dose. The recommended initial dose for patients age 6 years and older is 0.2 mg at bedtime. The dose may be titrated up to 0.6 mg to achieve the desired response. Fluid restriction should be observed, and fluid intake should be limited to a minimum from 1 hour before desmopressin administration, until the next morning, or at least 8 hours after administration. (See WARNINGS, PRECAUTIONS, Pediatric Use and Geriatric Use.)

HOW SUPPLIED

Strength	Size	NDC 68482-	Color	Markings
0.1 mg	Bottle of 30	275-30	White to off-white	
	Bottle of 90	275-90		
	Bottle of 100	275-01		
	Bottle of 1000	275-10		
0.2 mg	Bottle of 30	276-30	White to off-white	
	Bottle of 90	276-90		
	Bottle of 100	276-01		
	Bottle of 1000	276-10		

Desmopressin Acetate Tablets 0.1mg: White to off-white, oval, biconvex, uncoated tablets debossed "S" on one side and scored on the other side.

Desmopressin Acetate Tablets 0.2 mg: White to off-white, round, biconvex, uncoated tablets debossed "I" on one side and scored on the other side.

Store at 20° to 25°C (68° to 77°F) (see USP Controlled Room Temperature). Avoid exposure to excessive heat or light. This product should be dispensed in a container with a child-resistant cap.

Keep out of the reach of children.

Manufactured by:
 Glenmark Pharmaceuticals Ltd,
 Cochin-Bardes, Box 403 513, India

Manufactured for:

Glenmark Pharmaceuticals Inc., USA
 Mahwah, NJ 07430

Call center? 1 (888)721-7115
 www.glenmarkpharma.com/usa

March 2015

PE324251-1

		DATE: 04/03/2015	PANTONE SHADE NO: Black
PRODUCT NAME: LFLT DESMOPRESSIN ACETATE TAB US 6PI-AE	PKG. DEV.:	Item code, Version, Consistency of Design, copyright area, Pack size, Dimensions & Layout	<i>Shadhan 2/14/15</i>
ITEM CODE: PE32425 VERSION: 0315-1	RA:	Regulatory Text	<i>Shadhan 2/14/15</i>
PHARMACODE:	QA:	Quality Text	<i>Shadhan 2/14/15</i>
COUNTRY:	PRODUCTION:	Alphabetic Suitability	<i>Shadhan 2/14/15</i>
LOCATION: Goa	REMARKS:		
PACK: 40 GSM BIBLE PAPER			
ACTUAL SIZE: 200X280 MM			
SPECIFICATION: 32X32 MM			